

## TRACE ELEMENTS IN SEDIMENTS FROM PONDS IN PANTANAL, MATO GROSSO DO SUL, BRASIL

<sup>1</sup> Sandra R. Damatto, <sup>1</sup> Paulo S. C. Silva, <sup>1</sup> Barbara P. Mazzilli, <sup>2</sup> Arnaldo Y. Sakamoto, <sup>3</sup> Déborah I. T. Fávoro

<sup>1</sup> Centro de Metrologia das Radiações – Laboratório de Radiometria Ambiental - Instituto de Pesquisas Energéticas e Nucleares IPEN - Av. Prof. Lineu Prestes 2242, Cidade Universitária - 05508-000, São Paulo, BRASIL, [damatto@ipen.br](mailto:damatto@ipen.br)

<sup>2</sup> DCH/CPTL/Universidade Federal do Mato Grosso do Sul – Três Lagoas, Mato Grosso do Sul, BRASIL

<sup>3</sup> Centro do Reator de Pesquisas -Laboratório de Análise por Ativação Neutrônica - IPEN

### Abstract

Pantanal, in the southwest Brazil, is one of the world's largest freshwater wetland. This natural ecosystem has been affected due to urban contamination, irregular use of the land, tourism without control, excessive agricultural defensive utilization, etc. In order to verify possible changes in this environment, a study was established in Pantanal da Nhecolândia, Mato Grosso do Sul, Brazil. In 2006 two sediment cores were collected in two small ponds in Nhumirim farm, Salina do Meio and Salina da Ponta. The elements As, Ba, Br, Ce, Co, Cr, Cs, Eu, Fe, Hf, La, Lu, Na, Nd, Rb, Sb, Sc, Sm, Ta, Tb, Th, U, Yb, Zn and Zr were determined by instrumental neutron activation analysis technique – NAA. Their concentrations ranged from mg kg<sup>-1</sup> to %. For validation methodology the reference materials Buffalo River Sediment (NIST SRM 2704) and Soil-7 (IAEA) were analyzed. The majority of the elements and rare earths analyzed showed concentrations lower when compared with shale and earth crust values. Factorial analysis, mode R, Cluster analysis and principal component analysis were applied for the data interpretation.